

表 468 ETRF装着率(%) 細菌数測定の種類別 (ベッドサイドコンソール台数 1)

細菌数測定の 頻度	0 (装着せず)	10未満	10～	20～	30～	40～	50～	60～	70～	80～	90～	100 (全台装着)	合計	記載なし	総計	平均	標準偏差
なし	40	3	5	7		2	4	5	2	1	5	110	184		184	68.32	43.22
(%)	(21.7)	(1.6)	(2.7)	(3.8)		(1.1)	(2.2)	(2.7)	(1.1)	(0.5)	(2.7)	(59.8)	(100.0)				
毎日											1	12	13		13	99.84	0.58
(%)											(7.7)	(92.3)	(100.0)				
毎週	2	1	2	1		2	3	2	4	2	9	120	148		148	92.99	20.57
(%)	(1.4)	(0.7)	(1.4)	(0.7)		(1.4)	(2.0)	(1.4)	(2.7)	(1.4)	(6.1)	(81.1)	(100.0)				
隔週	1	2	3	1		1	2	4	3	3	8	201	229		229	95.34	17.09
(%)	(0.4)	(0.9)	(1.3)	(0.4)		(0.4)	(0.9)	(1.7)	(1.3)	(1.3)	(3.5)	(87.8)	(100.0)				
毎月	37	32	35	25	29	27	40	29	37	59	127	2,409	2,886		2,886	92.90	21.19
(%)	(1.3)	(1.1)	(1.2)	(0.9)	(1.0)	(0.9)	(1.4)	(1.0)	(1.3)	(2.0)	(4.4)	(83.5)	(100.0)				
年数回	11	1	5	6	2	2	3	9	6	8	15	286	354		354	91.05	23.99
(%)	(3.1)	(0.3)	(1.4)	(1.7)	(0.6)	(0.6)	(0.8)	(2.5)	(1.7)	(2.3)	(4.2)	(80.8)	(100.0)				
年1回	7	2	6	2	1	4	3	4	2	7	18	369	425		425	94.41	19.55
(%)	(1.6)	(0.5)	(1.4)	(0.5)	(0.2)	(0.9)	(0.7)	(0.9)	(0.5)	(1.6)	(4.2)	(86.8)	(100.0)				
合計	98	41	56	42	32	38	55	53	54	80	183	3,507	4,239		4,239	91.99	23.01
(%)	(2.3)	(1.0)	(1.3)	(1.0)	(0.8)	(0.9)	(1.3)	(1.3)	(1.3)	(1.9)	(4.3)	(82.7)	(100.0)				
不明	14		2	2	1	1	3		3	1		49	76	2	78	72.58	40.94
(%)	(18.4)		(2.6)	(2.6)	(1.3)	(1.3)	(3.9)		(3.9)	(1.3)		(64.5)	(100.0)				
記載なし												1	1		1	100.00	
(%)												(100.0)	(100.0)				
総計	112	41	58	44	33	39	58	53	57	81	183	3,557	4,316	2	4,318	91.65	23.57
(%)	(2.6)	(0.9)	(1.3)	(1.0)	(0.8)	(0.9)	(1.3)	(1.2)	(1.3)	(1.9)	(4.2)	(82.4)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
##C0034##
(C)Japanese Society for Dialysis Therapy
2016年調査