

表 563 脳出血の既往 原疾患別 (透析患者全体)

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	63,783	4,425	68,208	244	23,331	91,783
(%)	(93.5)	(6.5)	(100.0)			
慢性腎盂腎炎	2,052	140	2,192	8	696	2,896
(%)	(93.6)	(6.4)	(100.0)			
急速進行性糸球体腎炎	1,816	81	1,897	9	632	2,538
(%)	(95.7)	(4.3)	(100.0)			
妊娠腎/妊娠中毒症	1,056	80	1,136	5	342	1,483
(%)	(93.0)	(7.0)	(100.0)			
その他の分類不能の腎炎	988	54	1,042	4	408	1,454
(%)	(94.8)	(5.2)	(100.0)			
多発性嚢胞腎	7,720	1,112	8,832	36	2,603	11,471
(%)	(87.4)	(12.6)	(100.0)			
腎硬化症	22,130	1,769	23,899	115	7,636	31,650
(%)	(92.6)	(7.4)	(100.0)			
悪性高血圧	1,851	176	2,027	28	666	2,721
(%)	(91.3)	(8.7)	(100.0)			
糖尿病性腎症	86,158	5,299	91,457	439	31,761	123,657
(%)	(94.2)	(5.8)	(100.0)			
SLE腎炎	1,582	82	1,664	8	550	2,222
(%)	(95.1)	(4.9)	(100.0)			
アミロイド腎	310	11	321	2	115	438
(%)	(96.6)	(3.4)	(100.0)			
痛風腎	785	42	827	3	226	1,056
(%)	(94.9)	(5.1)	(100.0)			
先天性代謝異常による腎不全	207	10	217		64	281
(%)	(95.4)	(4.6)	(100.0)			
腎・尿路結核	112	10	122		46	168
(%)	(91.8)	(8.2)	(100.0)			
腎・尿路結石	426	33	459	4	115	578
(%)	(92.8)	(7.2)	(100.0)			
腎・尿路腫瘍	682	33	715	4	240	959
(%)	(95.4)	(4.6)	(100.0)			
閉塞性尿路障害	527	30	557		186	743
(%)	(94.6)	(5.4)	(100.0)			
骨髄腫	239	5	244	2	77	323
(%)	(98.0)	(2.0)	(100.0)			
腎形成不全	484	21	505	1	158	664
(%)	(95.8)	(4.2)	(100.0)			
不明	20,843	1,666	22,509	137	8,766	31,412
(%)	(92.6)	(7.4)	(100.0)			
再導入	1,563	73	1,636	5	533	2,174
(%)	(95.5)	(4.5)	(100.0)			
その他	5,659	398	6,057	37	2,344	8,438
(%)	(93.4)	(6.6)	(100.0)			
合計	220,973	15,550	236,523	1,091	81,495	319,109
(%)	(93.4)	(6.6)	(100.0)			
記載なし						
(%)						
総計	220,973	15,550	236,523	1,091	81,495	319,109
(%)	(93.4)	(6.6)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D0216##

(C)Japanese Society for Dialysis Therapy

2016年調査