

表 988 透析前クレアチニン濃度(mg/dL) 医療機関別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

医療機関	2未満	2~	4~	6~	8~	10~	12~	14~	16~	合計	記載なし	総計	平均	標準偏差
国公立大学		5	41	84	65	55	27	9	1	287	7	294	8.63	2.67
(%)		(1.7)	(14.3)	(29.3)	(22.6)	(19.2)	(9.4)	(3.1)	(0.3)	(100.0)				
私立大学		11	57	189	238	233	125	44	16	913	117	1,030	9.74	2.77
(%)		(1.2)	(6.2)	(20.7)	(26.1)	(25.5)	(13.7)	(4.8)	(1.8)	(100.0)				
国立病院・機構等		8	18	62	95	89	43	14	2	331	5	336	9.60	2.66
(%)		(2.4)	(5.4)	(18.7)	(28.7)	(26.9)	(13.0)	(4.2)	(0.6)	(100.0)				
県市町村立・国保	3	56	361	1,269	2,477	2,485	1,393	501	158	8,703	708	9,411	10.16	2.64
(%)	(0.0)	(0.6)	(4.1)	(14.6)	(28.5)	(28.6)	(16.0)	(5.8)	(1.8)	(100.0)				
JCHO(旧社会保険)等	2	25	93	287	524	477	311	96	25	1,840	27	1,867	9.98	2.69
(%)	(0.1)	(1.4)	(5.1)	(15.6)	(28.5)	(25.9)	(16.9)	(5.2)	(1.4)	(100.0)				
厚生連等	2	48	220	653	1,284	1,197	734	277	72	4,487	326	4,813	10.06	2.70
(%)	(0.0)	(1.1)	(4.9)	(14.6)	(28.6)	(26.7)	(16.4)	(6.2)	(1.6)	(100.0)				
その他公的	1	44	200	716	1,324	1,333	757	262	74	4,711	236	4,947	10.10	2.62
(%)	(0.0)	(0.9)	(4.2)	(15.2)	(28.1)	(28.3)	(16.1)	(5.6)	(1.6)	(100.0)				
私立総合		36	175	540	903	947	545	221	76	3,443	115	3,558	10.14	2.76
(%)		(1.0)	(5.1)	(15.7)	(26.2)	(27.5)	(15.8)	(6.4)	(2.2)	(100.0)				
私立	34	886	3,738	8,355	12,382	11,089	6,430	2,495	787	46,196	1,224	47,420	9.66	2.93
(%)	(0.1)	(1.9)	(8.1)	(18.1)	(26.8)	(24.0)	(13.9)	(5.4)	(1.7)	(100.0)				
私立診療所	28	511	2,939	11,333	23,308	24,277	15,119	6,077	2,028	85,620	1,837	87,457	10.41	2.68
(%)	(0.0)	(0.6)	(3.4)	(13.2)	(27.2)	(28.4)	(17.7)	(7.1)	(2.4)	(100.0)				
総計	70	1,630	7,842	23,488	42,600	42,182	25,484	9,996	3,239	156,531	4,602	161,133	10.14	2.78
(%)	(0.0)	(1.0)	(5.0)	(15.0)	(27.2)	(26.9)	(16.3)	(6.4)	(2.1)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

#D2714##

(C)Japanese Society for Dialysis Therapy

2016年調査