

表 993 透析前クレアチニン濃度(mg/dL) 医療機関別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

医療機関	2未満	2~	4~	6~	8~	10~	12~	14~	16~	合計	記載なし	総計	平均	標準偏差
国公立大学			7	9	7	11	8	1		43	2	45	9.24	2.75
(%)			(16.3)	(20.9)	(16.3)	(25.6)	(18.6)	(2.3)		(100.0)				
私立大学		2	4	31	60	59	31	17	4	208	4	212	10.31	2.52
(%)		(1.0)	(1.9)	(14.9)	(28.8)	(28.4)	(14.9)	(8.2)	(1.9)	(100.0)				
国立病院・機構等	1	9	7	2	9	9	5	3	1	46		46	8.45	4.02
(%)	(2.2)	(19.6)	(15.2)	(4.3)	(19.6)	(19.6)	(10.9)	(6.5)	(2.2)	(100.0)				
県市町村立・国保	1	8	61	262	571	593	402	129	55	2,082	130	2,212	10.48	2.61
(%)	(0.0)	(0.4)	(2.9)	(12.6)	(27.4)	(28.5)	(19.3)	(6.2)	(2.6)	(100.0)				
JCHO(旧社会保険)等		1	12	57	127	147	80	40	19	483	6	489	10.73	2.62
(%)		(0.2)	(2.5)	(11.8)	(26.3)	(30.4)	(16.6)	(8.3)	(3.9)	(100.0)				
厚生連等		5	19	136	332	386	229	82	34	1,223	68	1,291	10.67	2.72
(%)		(0.4)	(1.6)	(11.1)	(27.1)	(31.6)	(18.7)	(6.7)	(2.8)	(100.0)				
その他公的		6	29	109	250	270	180	60	20	924	57	981	10.46	2.59
(%)		(0.6)	(3.1)	(11.8)	(27.1)	(29.2)	(19.5)	(6.5)	(2.2)	(100.0)				
私立総合		12	40	154	298	318	218	86	28	1,154	83	1,237	10.41	2.75
(%)		(1.0)	(3.5)	(13.3)	(25.8)	(27.6)	(18.9)	(7.5)	(2.4)	(100.0)				
私立	9	131	623	2,038	3,857	4,260	2,637	1,074	369	14,998	346	15,344	10.37	2.77
(%)	(0.1)	(0.9)	(4.2)	(13.6)	(25.7)	(28.4)	(17.6)	(7.2)	(2.5)	(100.0)				
私立診療所	3	117	867	4,014	9,219	10,410	7,051	3,047	1,090	35,818	655	36,473	10.74	2.65
(%)	(0.0)	(0.3)	(2.4)	(11.2)	(25.7)	(29.1)	(19.7)	(8.5)	(3.0)	(100.0)				
総計	14	291	1,669	6,812	14,730	16,463	10,841	4,539	1,620	56,979	1,351	58,330	10.62	2.69
(%)	(0.0)	(0.5)	(2.9)	(12.0)	(25.9)	(28.9)	(19.0)	(8.0)	(2.8)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

#D2724##

(C)Japanese Society for Dialysis Therapy

2016年調査