

表 1089 透析前アルブミン濃度(g/dL) 医療機関別 (施設血液透析患者 透析回数週3回)

医療機関	1.5未満	1.5~	2.0~	2.5~	3.0~	3.5~	4.0~	4.5~	5.0~	合計	記載なし	総計	平均	標準偏差
国公立大学	2	18	63	99	105	105	24			416	16	432	3.05	0.64
(%)	(0.5)	(4.3)	(15.1)	(23.8)	(25.2)	(25.2)	(5.8)			(100.0)				
私立大学	5	30	96	162	326	551	164	9	1	1,344	213	1,557	3.37	0.60
(%)	(0.4)	(2.2)	(7.1)	(12.1)	(24.3)	(41.0)	(12.2)	(0.7)	(0.1)	(100.0)				
国立病院・機構等	3	9	23	56	121	175	57	3	1	448	8	456	3.38	0.59
(%)	(0.7)	(2.0)	(5.1)	(12.5)	(27.0)	(39.1)	(12.7)	(0.7)	(0.2)	(100.0)				
県市町村立・国保	20	97	324	983	3,455	5,201	1,445	66	9	11,600	923	12,523	3.47	0.48
(%)	(0.2)	(0.8)	(2.8)	(8.5)	(29.8)	(44.8)	(12.5)	(0.6)	(0.1)	(100.0)				
JCHO(旧社会保険)等	3	22	74	204	784	1,033	255	14	1	2,390	57	2,447	3.44	0.48
(%)	(0.1)	(0.9)	(3.1)	(8.5)	(32.8)	(43.2)	(10.7)	(0.6)	(0.0)	(100.0)				
厚生連等	5	47	140	497	1,822	2,653	712	51	22	5,949	453	6,402	3.48	0.48
(%)	(0.1)	(0.8)	(2.4)	(8.4)	(30.6)	(44.6)	(12.0)	(0.9)	(0.4)	(100.0)				
その他公的	14	50	190	543	1,895	2,751	738	31	4	6,216	327	6,543	3.46	0.49
(%)	(0.2)	(0.8)	(3.1)	(8.7)	(30.5)	(44.3)	(11.9)	(0.5)	(0.1)	(100.0)				
私立総合	3	25	111	348	1,340	2,175	633	32	2	4,669	189	4,858	3.51	0.46
(%)	(0.1)	(0.5)	(2.4)	(7.5)	(28.7)	(46.6)	(13.6)	(0.7)	(0.0)	(100.0)				
私立	75	429	1,819	5,832	17,733	25,741	7,714	440	43	59,826	2,258	62,084	3.46	0.49
(%)	(0.1)	(0.7)	(3.0)	(9.7)	(29.6)	(43.0)	(12.9)	(0.7)	(0.1)	(100.0)				
私立診療所	11	103	770	4,679	26,945	56,345	18,517	840	25	108,235	3,999	112,234	3.61	0.39
(%)	(0.0)	(0.1)	(0.7)	(4.3)	(24.9)	(52.1)	(17.1)	(0.8)	(0.0)	(100.0)				
総計	141	830	3,610	13,403	54,526	96,730	30,259	1,486	108	201,093	8,443	209,536	3.54	0.45
(%)	(0.1)	(0.4)	(1.8)	(6.7)	(27.1)	(48.1)	(15.0)	(0.7)	(0.1)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D3824##

(C)Japanese Society for Dialysis Therapy

2016年調査