

表 1097 アルブミン濃度(g/dL) 透析歴(年)別 (腹膜透析患者)

透析歴	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
2年未満	7	37	149	452	902	709	172	14	3	2,445	1,173	3,618	3.24	0.54
(%)	(0.3)	(1.5)	(6.1)	(18.5)	(36.9)	(29.0)	(7.0)	(0.6)	(0.1)	(100.0)				
2年～	6	24	98	351	757	611	124	4	1	1,976	1,008	2,984	3.25	0.50
(%)	(0.3)	(1.2)	(5.0)	(17.8)	(38.3)	(30.9)	(6.3)	(0.2)	(0.1)	(100.0)				
5年～	7	13	48	188	457	311	40	3		1,067	508	1,575	3.23	0.49
(%)	(0.7)	(1.2)	(4.5)	(17.6)	(42.8)	(29.1)	(3.7)	(0.3)		(100.0)				
10年～	2	4	12	37	111	38	9			213	76	289	3.15	0.49
(%)	(0.9)	(1.9)	(5.6)	(17.4)	(52.1)	(17.8)	(4.2)			(100.0)				
15年～		2	6	13	31	9	2			63	22	85	3.08	0.50
(%)		(3.2)	(9.5)	(20.6)	(49.2)	(14.3)	(3.2)			(100.0)				
20年～			2	5	5	7	1			20	8	28	3.22	0.54
(%)			(10.0)	(25.0)	(25.0)	(35.0)	(5.0)			(100.0)				
25年～			1		3	1				5	4	9	3.06	0.61
(%)			(20.0)		(60.0)	(20.0)				(100.0)				
30年～				2	1	1				4	2	6	3.15	0.51
(%)				(50.0)	(25.0)	(25.0)				(100.0)				
35年～						1				1		1	3.80	
(%)						(100.0)				(100.0)				
40年～				1	1					2		2	3.10	0.28
(%)				(50.0)	(50.0)					(100.0)				
合計	22	80	316	1,049	2,268	1,688	348	21	4	5,796	2,801	8,597	3.24	0.52
(%)	(0.4)	(1.4)	(5.5)	(18.1)	(39.1)	(29.1)	(6.0)	(0.4)	(0.1)	(100.0)				
不明			2	1	2	1				6	12	18	2.90	0.53
(%)			(33.3)	(16.7)	(33.3)	(16.7)				(100.0)				
記載なし														
(%)														
総計	22	80	318	1,050	2,270	1,689	348	21	4	5,802	2,813	8,615	3.24	0.52
(%)	(0.4)	(1.4)	(5.5)	(18.1)	(39.1)	(29.1)	(6.0)	(0.4)	(0.1)	(100.0)				
平均	3.68	3.14	3.00	3.10	3.35	2.90	2.39	1.57	0.75	3.08	2.91	3.03		
標準偏差	3.63	3.88	3.98	3.89	3.72	3.30	2.96	2.29	0.96	3.62	3.30	3.52		

数値下のかっこ内は行方向の合計に対する%です。

##D3842##

(C)Japanese Society for Dialysis Therapy

2016年調査