

表 1136 透析前補正カルシウム濃度(mg/dL) 透析歴(年)別 (施設血液透析患者 透析回数週3回)

透析歴	7.0未満	7.0~	7.5~	8.0~	8.4~	8.5~	9.0~	9.5~	10.0~	10.1~	10.5~	11.0~	11.5~	12.0~	合計	記載なし	総計	平均	標準偏差
2年未満	317	427	1,573	4,547	1,974	14,266	13,225	6,092	662	1,470	716	265	99	75	45,708	2,543	48,251	8.97	0.72
(%)	(0.7)	(0.9)	(3.4)	(9.9)	(4.3)	(31.2)	(28.9)	(13.3)	(1.4)	(3.2)	(1.6)	(0.6)	(0.2)	(0.2)	(100.0)				
2年~	220	306	1,205	3,512	1,590	13,879	16,173	9,193	1,060	2,368	1,073	365	116	89	51,149	2,213	53,362	9.12	0.71
(%)	(0.4)	(0.6)	(2.4)	(6.9)	(3.1)	(27.1)	(31.6)	(18.0)	(2.1)	(4.6)	(2.1)	(0.7)	(0.2)	(0.2)	(100.0)				
5年~	197	241	980	2,746	1,345	11,678	15,658	10,559	1,373	3,329	1,570	531	193	128	50,528	2,196	52,724	9.23	0.73
(%)	(0.4)	(0.5)	(1.9)	(5.4)	(2.7)	(23.1)	(31.0)	(20.9)	(2.7)	(6.6)	(3.1)	(1.1)	(0.4)	(0.3)	(100.0)				
10年~	92	118	478	1,280	597	5,196	7,455	5,793	784	2,153	1,089	364	144	65	25,608	1,099	26,707	9.32	0.76
(%)	(0.4)	(0.5)	(1.9)	(5.0)	(2.3)	(20.3)	(29.1)	(22.6)	(3.1)	(8.4)	(4.3)	(1.4)	(0.6)	(0.3)	(100.0)				
15年~	61	75	300	727	293	2,627	3,650	3,067	419	1,112	650	206	66	63	13,316	632	13,948	9.33	0.80
(%)	(0.5)	(0.6)	(2.3)	(5.5)	(2.2)	(19.7)	(27.4)	(23.0)	(3.1)	(8.4)	(4.9)	(1.5)	(0.5)	(0.5)	(100.0)				
20年~	36	61	166	409	178	1,474	1,926	1,513	236	545	346	90	37	29	7,046	305	7,351	9.29	0.81
(%)	(0.5)	(0.9)	(2.4)	(5.8)	(2.5)	(20.9)	(27.3)	(21.5)	(3.3)	(7.7)	(4.9)	(1.3)	(0.5)	(0.4)	(100.0)				
25年~	19	38	104	230	105	719	946	728	120	302	177	46	22	10	3,566	157	3,723	9.27	0.83
(%)	(0.5)	(1.1)	(2.9)	(6.4)	(2.9)	(20.2)	(26.5)	(20.4)	(3.4)	(8.5)	(5.0)	(1.3)	(0.6)	(0.3)	(100.0)				
30年~	5	28	68	139	41	399	473	409	53	149	74	24	17	4	1,883	86	1,969	9.24	0.83
(%)	(0.3)	(1.5)	(3.6)	(7.4)	(2.2)	(21.2)	(25.1)	(21.7)	(2.8)	(7.9)	(3.9)	(1.3)	(0.9)	(0.2)	(100.0)				
35年~	4	7	42	79	25	181	250	195	21	75	38	17	4	5	943	53	996	9.23	0.83
(%)	(0.4)	(0.7)	(4.5)	(8.4)	(2.7)	(19.2)	(26.5)	(20.7)	(2.2)	(8.0)	(4.0)	(1.8)	(0.4)	(0.5)	(100.0)				
40年~	3	2	6	23	10	63	103	69	11	18	17	11	3		339	14	353	9.30	0.83
(%)	(0.9)	(0.6)	(1.8)	(6.8)	(2.9)	(18.6)	(30.4)	(20.4)	(3.2)	(5.3)	(5.0)	(3.2)	(0.9)		(100.0)				
合計	954	1,303	4,922	13,692	6,158	50,482	59,859	37,618	4,739	11,521	5,750	1,919	701	468	200,086	9,298	209,384	9.16	0.75
(%)	(0.5)	(0.7)	(2.5)	(6.8)	(3.1)	(25.2)	(29.9)	(18.8)	(2.4)	(5.8)	(2.9)	(1.0)	(0.4)	(0.2)	(100.0)				
不明	1		2	8	3	19	26	27	3	4	12	2	2	4	113	39	152	9.55	1.27
(%)	(0.9)		(1.8)	(7.1)	(2.7)	(16.8)	(23.0)	(23.9)	(2.7)	(3.5)	(10.6)	(1.8)	(1.8)	(3.5)	(100.0)				
記載なし																			
(%)																			
総計	955	1,303	4,924	13,700	6,161	50,501	59,885	37,645	4,742	11,525	5,762	1,921	703	472	200,199	9,337	209,536	9.16	0.75
(%)	(0.5)	(0.7)	(2.5)	(6.8)	(3.1)	(25.2)	(29.9)	(18.8)	(2.4)	(5.8)	(2.9)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	6.23	6.86	6.43	5.89	5.78	5.99	6.73	7.89	8.52	8.91	9.52	9.19	9.60	9.13	6.96	6.64	6.94		
標準偏差	7.34	8.34	7.74	7.22	6.90	6.74	6.87	7.22	7.38	7.45	7.75	7.82	8.18	7.90	7.13	7.23	7.14		

数値下のカッコ内は行方向の合計に対する%です。

#D4032##

(C)Japanese Society for Dialysis Therapy

2016年調査