

表 1216 PTH値(pg/mL) 原疾患別(腹膜透析患者) PTH測定法:whole-PTH

原疾患	20未満	20~	35~	50~	75~	100~	125~	150~	180~	210~	240~	300~	350~	400~	500~	合計	記載なし	総計	平均	標準偏差	
慢性糸球体腎炎 (%)	12 (5.7)	11 (5.3)	20 (9.6)	34 (16.3)	24 (11.5)	24 (11.5)	20 (9.6)	11 (5.3)	11 (5.3)	12 (5.7)	10 (4.8)	9 (4.3)	4 (1.9)	4 (1.9)	3 (1.4)	209 (100.0)	12	221	136.56	120.18	
慢性腎孟腎炎 (%)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	1 (16.7)	6 (100.0)	6	6	84.83	48.08	
急速進行性糸球体腎炎 (%)					1 (33.3)			1 (33.3)		1 (33.3)						3 (100.0)	3	3	119.00	64.21	
妊娠腎/妊娠中毒症 (%)			1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	1 (25.0)	4 (100.0)	4	4	78.50	35.98	
その他の分類不能の腎炎 (%)																7 (100.0)	7	7	186.00	82.39	
多発性囊胞腎 (%)	2 (11.8)	2 (11.8)			2 (11.8)	4 (23.5)	1 (5.9)	1 (5.9)	2 (11.8)	1 (5.9)	1 (5.9)	1 (5.9)	1 (5.9)	1 (5.9)	1 (5.9)	17 (100.0)	2	19	110.06	86.44	
腎硬化症 (%)	2 (3.1)	5 (7.7)	6 (9.2)	10 (15.4)	8 (12.3)	4 (6.2)	6 (9.2)		4 (6.2)	7 (10.8)	3 (4.6)	3 (4.6)	1 (1.5)	1 (1.5)	2 (3.1)	4 (6.2)	65 (100.0)	3	68	162.89	181.60
悪性高血圧 (%)	2 (3.1)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	1 (3.3)	6 (100.0)	2	8	133.33	181.91	
糖尿病性腎症 (%)	16 (7.8)	20 (9.8)	20 (9.8)	30 (14.7)	20 (9.8)	29 (14.2)	20 (9.8)		13 (6.4)	10 (4.9)	7 (3.4)	9 (4.4)	5 (2.5)	3 (1.5)	1 (0.5)	1 (0.5)	204 (100.0)	11	215	112.69	90.82
SLE腎炎 (%)	1 (20.0)				1 (20.0)				1 (20.0)	1 (20.0)						5 (100.0)	5	5	113.00	76.53	
アミロイド腎 (%)	1 (100.0)															1 (100.0)	1	1	32.00		
痛風腎 (%)																3 (100.0)	3	3	158.00	103.81	
先天性代謝異常による腎不全 (%)	1 (100.0)															1 (100.0)	1	1	7.00		
腎・尿路結核 (%)																1 (100.0)					
腎・尿路結石 (%)		1 (20.0)	1 (20.0)	1 (20.0)												5 (100.0)	5	5	137.00	153.89	
腎・尿路腫瘍 (%)																					
閉塞性尿路障害 (%)			2 (100.0)													2 (100.0)	2	2	40.50	6.36	
骨髄腫 (%)																	1 (100.0)	1	1		
腎形成不全 (%)																3 (100.0)	3	3	200.33	177.08	
不明 (%)	4 (11.4)	3 (8.6)	2 (5.7)	6 (17.1)	7 (20.0)	4 (11.4)	2 (5.7)	1 (2.9)	3 (8.6)	2 (5.7)						1 (2.9)	35 (100.0)	7	42	115.03	141.29
再導入 (%)	1 (33.3)		1 (33.3)													1 (33.3)	3 (100.0)	3	3	221.33	338.81
その他 (%)	2 (7.7)	3 (11.5)	4 (15.4)	3 (11.5)	2 (7.7)	4 (15.4)		2 (7.7)	1 (3.8)		4 (15.4)	1 (3.8)				26 (100.0)	26	26	129.54	87.39	
合計 (%)	39 (6.4)	48 (7.9)	58 (9.6)	92 (15.2)	70 (11.6)	71 (11.7)	55 (9.1)		39 (6.4)	35 (5.8)	23 (3.8)	30 (5.0)	17 (2.8)	9 (1.5)	9 (1.5)	10 (1.7)	605 (100.0)	38	643	128.55	120.18
記載なし (%)																					
総計 (%)	39 (6.4)	48 (7.9)	58 (9.6)	92 (15.2)	70 (11.6)	71 (11.7)	55 (9.1)		39 (6.4)	35 (5.8)	23 (3.8)	30 (5.0)	17 (2.8)	9 (1.5)	9 (1.5)	10 (1.7)	605 (100.0)	38	643	128.55	120.18

数値下のカッコ内は行方向の合計に対する%です。

##D4283##

(C)Japanese Society for Dialysis Therapy

2016年調査