

表 1219 透析前ヘモグロビン濃度(g/dL) 治療方法(新分類)別 (透析患者全体)

治療方法 (新分類)	5未満	5~	6~	7~	8~	9~	10~	11~	12~	13~	14~	合計	記載なし	総計	平均	標準偏差	
施設血液透析	266	389	889	3,260	10,919	33,274	68,774	60,195	23,764	6,285	2,441	210,456	23,248	233,704	10.76	1.32	
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.2)	(15.8)	(32.7)	(28.6)	(11.3)	(3.0)	(1.2)	(100.0)					
Off-line HDF	4	7	20	74	234	683	1,298	1,212	496	145	44	4,217	420	4,637	10.77	1.34	
(%)	(0.1)	(0.2)	(0.5)	(1.8)	(5.5)	(16.2)	(30.8)	(28.7)	(11.8)	(3.4)	(1.0)	(100.0)					
On-line HDF	19	29	112	458	1,774	7,157	17,930	16,723	6,644	1,904	791	53,541	5,575	59,116	10.94	1.23	
(%)	(0.0)	(0.1)	(0.2)	(0.9)	(3.3)	(13.4)	(33.5)	(31.2)	(12.4)	(3.6)	(1.5)	(100.0)					
Push/Pull HDF				4	4	7	15	18	8	2	5	63	3	66	11.03	1.82	
(%)				(6.3)	(6.3)	(11.1)	(23.8)	(28.6)	(12.7)	(3.2)	(7.9)	(100.0)					
バイオフィルトレーション				4	17	24	69	64	33	5	3	219	33	252	10.82	1.32	
(%)				(1.8)	(7.8)	(11.0)	(31.5)	(29.2)	(15.1)	(2.3)	(1.4)	(100.0)					
IHDF	9	4	26	109	489	1,659	3,586	2,860	1,058	256	118	10,174	554	10,728	10.76	1.23	
(%)	(0.1)	(0.0)	(0.3)	(1.1)	(4.8)	(16.3)	(35.2)	(28.1)	(10.4)	(2.5)	(1.2)	(100.0)					
血液濾過						8	3	3	4	1		19		19	10.69	1.32	
(%)						(42.1)	(15.8)	(15.8)	(21.1)	(5.3)		(100.0)					
血液吸着透析 (リクセル等使用)	1		3	14	65	230	449	348	120	31	9	1,270	85	1,355	10.69	1.21	
(%)	(0.1)		(0.2)	(1.1)	(5.1)	(18.1)	(35.4)	(27.4)	(9.4)	(2.4)	(0.7)	(100.0)					
在宅血液透析		1	2	2	17	58	98	151	100	41	26	496	121	617	11.43	1.53	
(%)		(0.2)	(0.4)	(0.4)	(3.4)	(11.7)	(19.8)	(30.4)	(20.2)	(8.3)	(5.2)	(100.0)					
腹膜透析 (CAPD)	2	2	5	56	157	479	940	1,036	473	138	41	3,329	1,644	4,973	10.92	1.32	
(%)	(0.1)	(0.1)	(0.2)	(1.7)	(4.7)	(14.4)	(28.2)	(31.1)	(14.2)	(4.1)	(1.2)	(100.0)					
腹膜透析 (APD)	1	5	7	30	92	264	458	550	244	75	38	1,764	873	2,637	10.91	1.42	
(%)	(0.1)	(0.3)	(0.4)	(1.7)	(5.2)	(15.0)	(26.0)	(31.2)	(13.8)	(4.3)	(2.2)	(100.0)					
腹膜透析 (CCPD)				3	9	34	101	200	207	92	40	10	696	309	1,005	10.94	1.38
(%)				(0.4)	(1.3)	(4.9)	(14.5)	(28.7)	(29.7)	(13.2)	(5.7)	(1.4)	(100.0)				
合計	302	437	1,067	4,020	13,802	43,944	93,820	83,367	33,036	8,923	3,526	286,244	32,865	319,109	10.80	1.30	
(%)	(0.1)	(0.2)	(0.4)	(1.4)	(4.8)	(15.4)	(32.8)	(29.1)	(11.5)	(3.1)	(1.2)	(100.0)					
記載なし																	
(%)																	
総計	302	437	1,067	4,020	13,802	43,944	93,820	83,367	33,036	8,923	3,526	286,244	32,865	319,109	10.80	1.30	
(%)	(0.1)	(0.2)	(0.4)	(1.4)	(4.8)	(15.4)	(32.8)	(29.1)	(11.5)	(3.1)	(1.2)	(100.0)					

数値下の括弧内は行方向の合計に対する%です。

\*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

#D4311##

(C)Japanese Society for Dialysis Therapy

2016年調査