

表 1230 透析前ヘモグロビン濃度(g/dL) 医療機関別 (施設血液透析患者 透析回数週3回)

医療機関	5未満	5~	6~	7~	8~	9~	10~	11~	12~	13~	14~	合計	記載なし	総計	平均	標準偏差
国公立大学	1		4	31	71	91	91	80	31	9	3	412	20	432	10.05	1.54
(%)	(0.2)		(1.0)	(7.5)	(17.2)	(22.1)	(22.1)	(19.4)	(7.5)	(2.2)	(0.7)	(100.0)				
私立大学	5	10	18	52	130	236	385	334	136	32	11	1,349	208	1,557	10.41	1.53
(%)	(0.4)	(0.7)	(1.3)	(3.9)	(9.6)	(17.5)	(28.5)	(24.8)	(10.1)	(2.4)	(0.8)	(100.0)				
国立病院機構等	1	2	4	15	39	70	99	95	34	13	4	376	80	456	10.41	1.56
(%)	(0.3)	(0.5)	(1.1)	(4.0)	(10.4)	(18.6)	(26.3)	(25.3)	(9.0)	(3.5)	(1.1)	(100.0)				
県市町村立・国保	11	12	58	246	752	1,871	3,521	3,101	1,415	397	159	11,543	980	12,523	10.75	1.40
(%)	(0.1)	(0.1)	(0.5)	(2.1)	(6.5)	(16.2)	(30.5)	(26.9)	(12.3)	(3.4)	(1.4)	(100.0)				
JCHO(旧社会保険)等	1	2	11	50	152	448	700	629	293	81	31	2,398	49	2,447	10.71	1.37
(%)	(0.0)	(0.1)	(0.5)	(2.1)	(6.3)	(18.7)	(29.2)	(26.2)	(12.2)	(3.4)	(1.3)	(100.0)				
厚生連等	3	7	46	102	314	920	1,780	1,774	758	237	99	6,040	362	6,402	10.84	1.39
(%)	(0.0)	(0.1)	(0.8)	(1.7)	(5.2)	(15.2)	(29.5)	(29.4)	(12.5)	(3.9)	(1.6)	(100.0)				
その他公的	8	5	38	122	352	912	1,805	1,755	838	248	107	6,190	353	6,543	10.85	1.43
(%)	(0.1)	(0.1)	(0.6)	(2.0)	(5.7)	(14.7)	(29.2)	(28.4)	(13.5)	(4.0)	(1.7)	(100.0)				
私立総合	3	7	20	89	248	690	1,498	1,293	566	174	73	4,661	197	4,858	10.81	1.37
(%)	(0.1)	(0.2)	(0.4)	(1.9)	(5.3)	(14.8)	(32.1)	(27.7)	(12.1)	(3.7)	(1.6)	(100.0)				
私立	84	150	335	1,158	3,667	9,968	18,581	16,370	7,046	1,976	727	60,062	2,022	62,084	10.72	1.38
(%)	(0.1)	(0.2)	(0.6)	(1.9)	(6.1)	(16.6)	(30.9)	(27.3)	(11.7)	(3.3)	(1.2)	(100.0)				
私立診療所	141	179	284	1,121	4,571	16,511	37,571	32,526	11,746	2,872	1,127	108,649	3,585	112,234	10.80	1.24
(%)	(0.1)	(0.2)	(0.3)	(1.0)	(4.2)	(15.2)	(34.6)	(29.9)	(10.8)	(2.6)	(1.0)	(100.0)				
総計	258	374	818	2,986	10,296	31,717	66,031	57,957	22,863	6,039	2,341	201,680	7,856	209,536	10.77	1.31
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.1)	(15.7)	(32.7)	(28.7)	(11.3)	(3.0)	(1.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4324##

(C)Japanese Society for Dialysis Therapy

2016年調査