

表 1247 総コレステロール濃度(mg/dL) 原疾患別 (透析患者全体)

原疾患	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	1,855	5,659	12,038	15,566	14,156	9,960	5,314	2,507	1,027	654	68,736	23,047	91,783	161.28	35.97
(%)	(2.7)	(8.2)	(17.5)	(22.6)	(20.6)	(14.5)	(7.7)	(3.6)	(1.5)	(1.0)	(100.0)				
慢性腎盂腎炎	56	161	350	472	480	331	208	86	43	25	2,212	684	2,896	164.07	36.61
(%)	(2.5)	(7.3)	(15.8)	(21.3)	(21.7)	(15.0)	(9.4)	(3.9)	(1.9)	(1.1)	(100.0)				
急速進行性糸球体腎炎	33	109	254	341	395	269	234	116	64	65	1,880	658	2,538	173.92	42.24
(%)	(1.8)	(5.8)	(13.5)	(18.1)	(21.0)	(14.3)	(12.4)	(6.2)	(3.4)	(3.5)	(100.0)				
妊娠腎/妊娠中毒症	11	54	146	193	246	215	135	84	31	29	1,144	339	1,483	175.20	38.39
(%)	(1.0)	(4.7)	(12.8)	(16.9)	(21.5)	(18.8)	(11.8)	(7.3)	(2.7)	(2.5)	(100.0)				
その他の分類不能の腎炎	29	93	186	241	236	149	105	44	19	10	1,112	342	1,454	162.49	37.19
(%)	(2.6)	(8.4)	(16.7)	(21.7)	(21.2)	(13.4)	(9.4)	(4.0)	(1.7)	(0.9)	(100.0)				
多発性嚢胞腎	178	670	1,475	1,980	1,891	1,299	723	302	98	63	8,679	2,792	11,471	161.96	34.53
(%)	(2.1)	(7.7)	(17.0)	(22.8)	(21.8)	(15.0)	(8.3)	(3.5)	(1.1)	(0.7)	(100.0)				
腎硬化症	707	2,340	4,733	5,698	4,670	2,942	1,499	673	262	197	23,721	7,929	31,650	156.88	35.01
(%)	(3.0)	(9.9)	(20.0)	(24.0)	(19.7)	(12.4)	(6.3)	(2.8)	(1.1)	(0.8)	(100.0)				
悪性高血圧	38	151	358	436	430	294	147	79	33	22	1,988	733	2,721	162.62	35.35
(%)	(1.9)	(7.6)	(18.0)	(21.9)	(21.6)	(14.8)	(7.4)	(4.0)	(1.7)	(1.1)	(100.0)				
糖尿病性腎症	4,715	12,445	20,389	21,122	15,847	9,405	4,667	2,044	874	639	92,147	31,510	123,657	150.70	35.63
(%)	(5.1)	(13.5)	(22.1)	(22.9)	(17.2)	(10.2)	(5.1)	(2.2)	(0.9)	(0.7)	(100.0)				
SLE腎炎	31	106	229	339	346	246	185	117	50	33	1,682	540	2,222	170.76	39.47
(%)	(1.8)	(6.3)	(13.6)	(20.2)	(20.6)	(14.6)	(11.0)	(7.0)	(3.0)	(2.0)	(100.0)				
アミロイド腎	7	26	64	62	53	50	27	10	6	6	311	127	438	162.32	38.80
(%)	(2.3)	(8.4)	(20.6)	(19.9)	(17.0)	(16.1)	(8.7)	(3.2)	(1.9)	(1.9)	(100.0)				
痛風腎	34	73	159	182	174	95	50	13	13	10	803	253	1,056	156.64	35.45
(%)	(4.2)	(9.1)	(19.8)	(22.7)	(21.7)	(11.8)	(6.2)	(1.6)	(1.6)	(1.2)	(100.0)				
先天性代謝異常による腎不全	7	14	33	54	34	23	22	5	4	8	204	77	281	164.01	42.13
(%)	(3.4)	(6.9)	(16.2)	(26.5)	(16.7)	(11.3)	(10.8)	(2.5)	(2.0)	(3.9)	(100.0)				
腎・尿路結核	3	9	23	29	20	7	17	6	1	1	116	52	168	161.74	36.93
(%)	(2.6)	(7.8)	(19.8)	(25.0)	(17.2)	(6.0)	(14.7)	(5.2)	(0.9)	(0.9)	(100.0)				
腎・尿路結石	18	34	81	101	92	53	34	12	1	4	430	148	578	157.84	34.83
(%)	(4.2)	(7.9)	(18.8)	(23.5)	(21.4)	(12.3)	(7.9)	(2.8)	(0.2)	(0.9)	(100.0)				
腎・尿路腫瘍	26	77	160	169	135	71	41	22	6	3	710	249	959	153.30	33.96
(%)	(3.7)	(10.8)	(22.5)	(23.8)	(19.0)	(10.0)	(5.8)	(3.1)	(0.8)	(0.4)	(100.0)				
閉塞性尿路障害	6	40	101	121	117	81	56	25	4	4	555	188	743	163.43	34.15
(%)	(1.1)	(7.2)	(18.2)	(21.8)	(21.1)	(14.6)	(10.1)	(4.5)	(0.7)	(0.7)	(100.0)				
骨髄腫	8	21	40	40	35	30	21	11	7	7	220	103	323	165.75	45.95
(%)	(3.6)	(9.5)	(18.2)	(18.2)	(15.9)	(13.6)	(9.5)	(5.0)	(3.2)	(3.2)	(100.0)				
腎形成不全	9	29	89	98	102	84	46	15	9	7	488	176	664	165.46	37.43
(%)	(1.8)	(5.9)	(18.2)	(20.1)	(20.9)	(17.2)	(9.4)	(3.1)	(1.8)	(1.4)	(100.0)				
不明	701	2,205	4,157	5,207	4,446	2,929	1,678	729	288	219	22,559	8,853	31,412	158.63	36.34
(%)	(3.1)	(9.8)	(18.4)	(23.1)	(19.7)	(13.0)	(7.4)	(3.2)	(1.3)	(1.0)	(100.0)				
再導入	41	127	273	374	344	224	135	78	28	26	1,650	524	2,174	163.85	37.64
(%)	(2.5)	(7.7)	(16.5)	(22.7)	(20.8)	(13.6)	(8.2)	(4.7)	(1.7)	(1.6)	(100.0)				
その他	245	556	1,076	1,312	1,223	812	476	206	110	77	6,093	2,345	8,438	160.05	38.53
(%)	(4.0)	(9.1)	(17.7)	(21.5)	(20.1)	(13.3)	(7.8)	(3.4)	(1.8)	(1.3)	(100.0)				
合計	8,758	24,999	46,414	54,137	45,472	29,569	15,820	7,184	2,978	2,109	237,440	81,669	319,109	156.74	36.32
(%)	(3.7)	(10.5)	(19.5)	(22.8)	(19.2)	(12.5)	(6.7)	(3.0)	(1.3)	(0.9)	(100.0)				
記載なし															
(%)															
総計	8,758	24,999	46,414	54,137	45,472	29,569	15,820	7,184	2,978	2,109	237,440	81,669	319,109	156.74	36.32
(%)	(3.7)	(10.5)	(19.5)	(22.8)	(19.2)	(12.5)	(6.7)	(3.0)	(1.3)	(0.9)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4416##

(C)Japanese Society for Dialysis Therapy

2016年調査