

表 1262 総コレステロール濃度(mg/dL) 原疾患別 (腹膜透析患者)

原疾患	100未満	100~	120~	140~	160~	180~	200~	220~	240~	260~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	22	54	143	239	301	299	173	107	60	57	1,455	1,211	2,666	180.06	40.76
(%)	(1.5)	(3.7)	(9.8)	(16.4)	(20.7)	(20.5)	(11.9)	(7.4)	(4.1)	(3.9)	(100.0)				
慢性腎盂腎炎	1	4	6	1	7	15	12	6	1		53	44	97	181.34	38.18
(%)	(1.9)	(7.5)	(11.3)	(1.9)	(13.2)	(28.3)	(22.6)	(11.3)	(1.9)		(100.0)				
急速進行性糸球体腎炎		3	3	7	9	4	1	5	4	36	25	61	203.50	53.75	
(%)		(8.3)	(8.3)	(19.4)	(25.0)	(11.1)	(2.8)	(13.9)	(11.1)	(100.0)					
妊娠腎/妊娠中毒症		2		3	3	2	5	1	1	17	7	24	203.18	46.25	
(%)		(11.8)		(17.6)	(17.6)	(11.8)	(29.4)	(5.9)	(5.9)	(100.0)					
その他の分類不能の腎炎	1	3	5	10	10	7	3	3	3	45	30	75	191.18	43.64	
(%)	(2.2)	(6.7)	(11.1)	(22.2)	(22.2)	(15.6)	(6.7)	(6.7)	(6.7)	(100.0)					
多発性囊胞腎	3	7	24	30	24	21	17	6	6	138	134	272	188.49	40.44	
(%)	(2.2)	(5.1)	(17.4)	(21.7)	(17.4)	(15.2)	(12.3)	(4.3)	(4.3)	(100.0)					
腎硬化症	22	54	110	127	100	62	51	15	19	560	546	1,106	179.34	39.79	
(%)	(3.9)	(9.6)	(19.6)	(22.7)	(17.9)	(11.1)	(9.1)	(2.7)	(3.4)	(100.0)					
悪性高血圧	2	5	10	16	9	4	4	4	2	56	50	106	180.54	38.53	
(%)	(3.6)	(8.9)	(17.9)	(28.6)	(16.1)	(7.1)	(7.1)	(7.1)	(3.6)	(100.0)					
糖尿病性腎症	30	111	221	283	306	210	126	81	25	36	1,429	1,317	2,746	167.10	39.95
(%)	(2.1)	(7.8)	(15.5)	(19.8)	(21.4)	(14.7)	(8.8)	(5.7)	(1.7)	(2.5)	(100.0)				
SLE腎炎	1		1	2	9	2	3	6	4		28	27	55	193.86	42.24
(%)	(3.6)		(3.6)	(7.1)	(32.1)	(7.1)	(10.7)	(21.4)	(14.3)		(100.0)				
アミロイド腎		2	2	1						1	6	10	16	161.83	53.52
(%)		(33.3)	(33.3)	(16.7)						(16.7)	(100.0)				
痛風腎	1	1	1	3	3	1	3		1	14	9	23	167.21	41.40	
(%)	(7.1)	(7.1)	(7.1)	(21.4)	(21.4)	(7.1)	(21.4)		(7.1)	(100.0)					
先天性代謝異常による腎不全	1			3	2	1	3			10	7	17	171.00	34.89	
(%)	(10.0)			(30.0)	(20.0)	(10.0)	(30.0)			(100.0)					
腎・尿路結核											1	1			
(%)															
腎・尿路結石	1	1	1	1			3	4		1	12	9	21	197.75	45.01
(%)	(8.3)	(8.3)	(8.3)	(8.3)	(8.3)		(25.0)	(33.3)		(8.3)	(100.0)				
腎・尿路腫瘍		1	2			1				4	8	12	154.00	19.36	
(%)		(25.0)	(50.0)			(25.0)				(100.0)					
閉塞性尿路障害	3	1		4	2	1				11	9	20	177.55	34.90	
(%)	(27.3)	(9.1)		(36.4)	(18.2)	(9.1)				(100.0)					
骨髄腫					2	1				3	4	7	204.67	26.50	
(%)					(66.7)	(33.3)				(100.0)					
腎形成不全	1		4	2	7	7	6	4	2	1	34	31	65	186.38	41.24
(%)	(2.9)		(11.8)	(5.9)	(20.6)	(20.6)	(17.6)	(11.8)	(5.9)	(2.9)	(100.0)				
不明	3	13	33	88	88	83	70	32	21	18	449	440	889	182.67	40.86
(%)	(0.7)	(2.9)	(7.3)	(19.6)	(19.6)	(18.5)	(15.6)	(7.1)	(4.7)	(4.0)	(100.0)				
再導入	2	3	12	12	6	7	4	3	1	1	39	26	65	174.90	37.92
(%)	(5.1)	(7.7)	(30.8)	(15.4)	(17.9)	(10.3)	(7.7)	(2.6)	(2.6)	(100.0)					
その他	3	12	24	25	26	21	8	5	5	129	142	271	184.67	45.09	
(%)	(2.3)	(9.3)	(18.6)	(19.4)	(20.2)	(16.3)	(6.2)	(3.9)	(3.9)	(100.0)					
合計	60	217	505	815	949	813	526	334	154	155	4,528	4,087	8,615	176.97	41.25
(%)	(1.3)	(4.8)	(11.2)	(18.0)	(21.0)	(18.0)	(11.6)	(7.4)	(3.4)	(3.4)	(100.0)				
記載なし															
(%)															
総計	60	217	505	815	949	813	526	334	154	155	4,528	4,087	8,615	176.97	41.25
(%)	(1.3)	(4.8)	(11.2)	(18.0)	(21.0)	(18.0)	(11.6)	(7.4)	(3.4)	(3.4)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4443##

(C)Japanese Society for Dialysis Therapy

2016年調査