

表 1266 HDL-C濃度(mg/dL) 併用の状況別 (透析患者全体)

併用の状況	20未満	20 ~	30 ~	40 ~	50 ~	60 ~	70 ~	80 ~	90 ~	100 ~	合計	記載なし	総計	平均	標準偏差
HD(F)単独	814	17,081	51,588	58,519	43,177	25,361	12,635	5,763	2,567	2,120	219,625	90,513	310,138	48.80	16.46
(%)	(0.4)	(7.8)	(23.5)	(26.6)	(19.7)	(11.5)	(5.8)	(2.6)	(1.2)	(1.0)	(100.0)				
PD単独	10	163	682	784	632	406	209	123	62	48	3,119	3,812	6,931	51.54	18.38
(%)	(0.3)	(5.2)	(21.9)	(25.1)	(20.3)	(13.0)	(6.7)	(3.9)	(2.0)	(1.5)	(100.0)				
HD(F)単独.PDカテあり		9	31	29	29	9	8	7	2		124	53	177	49.48	16.44
(%)		(7.3)	(25.0)	(23.4)	(23.4)	(7.3)	(6.5)	(5.6)	(1.6)		(100.0)				
PD.HD併用週1HD(F)	5	119	219	224	133	98	33	23	5	8	867	629	1,496	46.07	16.85
(%)	(0.6)	(13.7)	(25.3)	(25.8)	(15.3)	(11.3)	(3.8)	(2.7)	(0.6)	(0.9)	(100.0)				
PD.HD併用週2HD(F)		8	22	16	22	14	4	2			88	77	165	48.01	14.72
(%)		(9.1)	(25.0)	(18.2)	(25.0)	(15.9)	(4.5)	(2.3)			(100.0)				
PD.HD併用週3HD(F)		3	5	6	2	1	4				21	16	37	48.05	17.64
(%)		(14.3)	(23.8)	(28.6)	(9.5)	(4.8)	(19.0)				(100.0)				
PD.HD併用週4HD(F)											1	1			
(%)															
PD.HD併用その他		2	10	15	9	5	1	1			43	20	63	47.14	13.13
(%)		(4.7)	(23.3)	(34.9)	(20.9)	(11.6)	(2.3)	(2.3)			(100.0)				
合計	829	17,385	52,557	59,593	44,004	25,894	12,894	5,919	2,636	2,176	223,887	95,121	319,008	48.83	16.49
(%)	(0.4)	(7.8)	(23.5)	(26.6)	(19.7)	(11.6)	(5.8)	(2.6)	(1.2)	(1.0)	(100.0)				
不明				2		1		1		1	5	17	22	72.40	34.20
(%)				(40.0)		(20.0)		(20.0)		(20.0)	(100.0)				
記載なし											79	79			
(%)															
総計	829	17,385	52,557	59,595	44,004	25,895	12,894	5,920	2,636	2,177	223,892	95,217	319,109	48.83	16.49
(%)	(0.4)	(7.8)	(23.5)	(26.6)	(19.7)	(11.6)	(5.8)	(2.6)	(1.2)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4512##

(C)Japanese Society for Dialysis Therapy  
2016年調査